The separate African winter quarters of Pied Flycatcher Ficedula hypoleuca and Collared Flycatcher F. albicollis

Robert J. Dowsett

Le Gobemouche noir Ficedula bypoleuca et la Gobemouche à collier F. albicollis ont des quartiers d'hiver africains bien séparés. Goodenough et al. (2009) ont écrit que le Gobemouche noir Ficedula hypoleuca et son proche cousin le Gobemouche à collier F. albicollis hivernent en sympatrie dans des pays comme le Tchad et le Nigeria, mais c'est incorrect. Le Gobemouche noir hiverne principalement dans les forêts claires soudaniennes de la Sierra Leone au Cameroun, tandis que le Gobemouche à collier hiverne dans les forêts claires de la région zambézienne au sud de l'Equateur.

Summary. The claim that Pied Flycatcher *Ficedula hypoleuca* and its close relative the Collared Flycatcher *F. albicollis* 'winter sympatrically in countries such as Chad and Nigeria' (Goodenough *et al.* 2009) is not true. Pied Flycatcher winters mainly in the Sudanian woodlands from Cameroon west to Sierra Leone, but Collared Flycatcher does so in the Zambezian woodlands south of the equator.

In their discussion of factors relevant to the conservation of Pied Flycatchers *Ficedula* hypoleuca, Goodenough et al. (2009) claim that this species and its close relative the Collared Flycatcher *F. albicollis* 'winter sympatrically in countries such as Chad and Nigeria (Dowsett 1993, Elgood et al. 1994)'. This is untrue, and is not what was shown by Dowsett (1993).

A literature search has produced many specimen and correctly identified sight records of *Ficedula* flycatchers in sub-Saharan Africa. To these have been added my own observations from several countries in which these two species occur in winter or on passage, from western Mali to northern Cameroon for Pied Flycatchers, and north-east Nigeria to southern Zambia for Collared Flycatchers.

Winter distribution of Pied and Collared Flycatchers

Goodenough *et al.* (2009) were perhaps influenced by the fact that Elgood *et al.* (1994) wrote of Collared Flycatcher in Nigeria 'uncommon... Sep-Mar', which implies a presence throughout that period. This echoed Elgood *et al.* (1966), who considered that it is 'probably a regular winter visitor'. However, the few records of Collared Flycatcher in Nigeria all fall between 13 September and 1 November, except for one bird which spent the period 23 February-25 March at Vom (Smith 1966). There is a handful of records from neighbouring Chad (11 September-19 October and 11-15 April: Salvan 1968). For both Nigeria and Chad, Dowsett (1993) made it clear that the species occurs only on passage.

Pied Flycatchers too are known only on passage through Chad (7 September to the first half of October and 29 March-10 April: Salvan 1968, Newby 1980), which is at the eastern limit of the species' migration. But they do winter in considerable numbers in Nigeria. Elgood et al. (1994) reported them between October and April, but extreme dates are in fact 17 September (Malamfatori, Lake Chad: Dowsett 1969) and 23 May (Vom: Smith 1966). Pied Flycatchers winter south in Nigeria from 11°N (Falgore, January: Wilkinson & Beecroft 1985; Zaria, 27 February: Fry 1965), mostly between 07°N and 10°N. There are few records from the coastal zone of Nigeria (e.g. Lagos: Forrester 1971, Gee & Heigham 1977). In its overall wintering area (essentially Cameroon west to Mali, south to Sierra Leone: Serle 1949; pers. obs.) the species frequents especially the Sudanian woodlands, less often clearings or the edge of forest. In the Sahel it occurs only on passage, and then there is a preponderance of autumn (southbound passage) records at several localities; e.g. at Malamfatori, at least 14 between 17 September and 9 October (Dowsett 1969), but only three on northbound passage (early April to 3 May: Dowsett 1968, Sharland 1969). In contrast, with the drying out of the Lake Chad area, Gustafsson et al. (2003) in the period 1997-2000 had only a single autumn record at Malamfatori (8 September), and none at all in spring. Near Jos (2001-09), M. Stevens (in




Figure 1. Records (mapped by degree square) of Pied Flycatchers *Ficedula hypoleuca* (black squares), Collared Flycatchers *F. albicollis* (open squares) and Semi-collared Flycatchers *F. semitorquata* (black dots). Localities at which two species occur are denoted by 'pied' symbols.

Distribution (par carrés d'un degré de latitude-longitude) des Gobemouches noirs *Ficedula hypoleuca* (carrés noirs), à collier *F. albicollis* (carrés creux) et à demi-collier *F. semitorquata* (points noirs). Les localités où se rencontrent deux espèces sont indiquées par un symbole noir et blanc.

litt. 2009) ringed 100 in October compared to 45 in March–April.

In contrast to Pied Flycatchers, Collared Flycatchers winter south of the equator, in the Zambezian woodlands, centred on Zambia and Malaŵi in particular (Moreau 1972, Dowsett-Lemaire & Dowsett 2006, Dowsett *et al.* 2008). Fig. 1 shows the distribution of these two species, and the separation of the centres of distribution is clear, with the Central African rain forest largely avoided. Records of Semi-collared Flycatchers *F. semitorquata* are also mapped; as they are difficult to separate, some records might have to be reassessed, but it does seem that their winter distribution is near the equator, centred on Uganda and western Kenya (e.g. Pearson 1998, and references therein).

Future workers are likely to be misled by the published statement in such an authoritative journal as *Ibis*, but the editor declined to publish this clarification there.

Acknowledgements

I thank Peter Jones, David Pearson and Matt Stevens for their helpful comments on the manuscript.

References

- Dowsett, R. J. 1968. Migrants at Malam'fatori, Lake Chad, spring 1968. *Bull. Nigerian Orn. Soc.* 5: 53–56.
- Dowsett, R. J. 1969. Migrants at Malam'fatori, Lake Chad, autumn 1968. *Bull. Nigerian Orn. Soc.* 6: 39–45.
- Dowsett, R. J. 1993. Afrotropical avifaunas: annotated country lists. Nigeria and Chad. *Tauraco Res. Rep.* 5: 69–77 & 103–108.
- Dowsett, R. J., Aspinwall, D. R. & Dowsett-Lemaire, F. 2008. *The Birds of Zambia. An Atlas and Handbook.* Liège: Tauraco Press & Aves.

African winter quarters Pied and Collared Flycatchers: Dowsett

- Dowsett-Lemaire, F. & Dowsett, R. J. 2006. *The Birds* of Malawi. An Atlas and Handbook. Liège: Tauraco Press & Aves.
- Elgood, J. H., Sharland, R. E. & Ward, P. 1966. Palaearctic migrants in Nigeria. *Ibis* 108: 84–116.
- Elgood, J. H., Heigham, J. B., Moore, A. M., Nason, A. M., Sharland, R. E. & Skinner, N. J. 1994. *The Birds of Nigeria: An Annotated Check-list*. BOU Check-list No. 4. Second edn. Tring: British Ornithologists' Union.
- Forrester, M. F. 1971. The birds of Ikoyi, Lagos. Bull. Nigerian Orn. Soc. 8: 13–21.
- Fry, C. H. 1965. The birds of Zaria. IV—Residents, vagrants and check-list (Passerines). *Bull. Nigerian* Orn. Soc. 2: 91–102.
- Gee, J. & Heigham, J. 1977. Birds of Lagos, Nigeria. [Part 2]. Bull. Nigerian Orn. Soc. 13: 103–132.
- Goodenough, A. E., Elliot, S. L. & Hart, A. G. 2009. The challenges of conservation for declining migrants: are reserve-based initiatives during the breeding season appropriate for the Pied Flycatcher *Ficedula hypoleuca? Ibis* 151: 429–439.
- Gustafsson, R., Hjort, C., Ottosson, U. & Hall, P. 2003. Birds at Lake Chad and in the Sahel of NE Nigeria 1997–2000. Ottenby: Lake Chad Bird Migration Project.

- Moreau, R. E. 1972. *The Palaearctic–African Bird Migration System*. London, UK: Academic Press.
- Newby, J. E. 1980. The birds of the Oadi Rime— Ouadi Achim Faunal Reserve, a contribution to the study of the Chadian avifauna. [Part 2.] *Malimbus* 2: 29–50.
- Pearson, D. J. 1998. Pied Flycatcher *Ficedula hypoleuca* at Kakamega Forest—a readmission to the East African list. *Scopus* 20: 46–48.
- Salvan, J. 1968. Contribution à l'étude des oiseaux du Tchad. [Part 4.] *Oiseau & R.F.O.* 38: 249–273.
- Serle, W. 1949. Notes on the birds of Sierra Leone. [Part 3]. Ostrich 20: 70–85.
- Sharland, R. E. 1969. Mallam Fatori, spring 1969. Bull. Nigerian Orn. Soc. 6: 98–101.
- Smith, V. W. 1966. Autumn and spring weights of some Palaearctic migrants in central Nigeria. *Ibis* 108: 492–512.
- Wilkinson, R. & Beecroft, R. 1985. Birds in Falgore Game Reserve, Nigeria. *Malimbus* 7: 63–72.

Le Pouget, 30440 Sumène, France. E-mail: Dowsett@ aol.com

Received 15 September 2009; revision accepted 20 October 2009.